

# Accepted Manuscript

Title: Could titin have a role in strain-induced injuries?

Author: Craig Perrin, Kazunori Nosaka, James Steele

PII: S2095-2546(17)30031-5

DOI: <http://dx.doi.org/doi: 10.1016/j.jshs.2017.03.003>

Reference: JSHS 376

To appear in: *Journal of Sport and Health Science*

Received date: 21-2-2017

Accepted date: 27-2-2017

Please cite this article as: Craig Perrin, Kazunori Nosaka, James Steele, Could titin have a role in strain-induced injuries?, *Journal of Sport and Health Science* (2017), <http://dx.doi.org/doi: 10.1016/j.jshs.2017.03.003>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



## Commentary

### Could titin have a role in strain-induced injuries?

Craig Perrin <sup>a\*</sup>, Kazunori Nosaka <sup>b</sup>, James Steele <sup>a</sup>

<sup>a</sup> School of Sport, Health and Social Sciences, Southampton Solent University Ringgold standard institution, Southampton, SO14 0YN, UK

<sup>b</sup> School of Exercise and Health Sciences, Edith Cowan University, Joondalup, WA 6027, Australia

\*Corresponding author: Craig Perrin

Email: [Operrc41@solent.ac.uk](mailto:Operrc41@solent.ac.uk)

Running head: Titin and strain-induced injuries

Received 21 February 2017; accepted 27 February 2017

It has been known for some time that strain injuries occur through the excessive lengthening of a muscle.<sup>1-3</sup> However, the precise mechanism for strain injury remains elusive and has been the subject of recent debate in the *Journal of Sport and Health Science*,<sup>4-9</sup> but it seems sensible to consider mechanisms at the level of the muscle fiber. It was previously believed that damage was the result of non-uniform stretching of sarcomeres on the descending limb of the length — tension curve, with the weakest sarcomeres undergoing the greatest deformation and ultimately damaging the myofibril.<sup>10</sup> This non-uniform lengthening was also believed to explain the residual force enhancement observed during lengthening contractions, as the rapid lengthening of some sarcomeres gives rise to passive tension whilst allowing the remaining sarcomeres to operate closer to their optimal length.<sup>11</sup> Whilst it has been shown sarcomeres do lengthen non-uniformly, it does not appear to be dependent on the strength of the sarcomere.<sup>12</sup> Furthermore, the increase in sarcomere length non-uniformity after active stretching is not correlated to the residual increase in force ( $r = -0.309$ ).<sup>12</sup> Although this does not disprove the notion that sarcomere lengthening is responsible for muscle damage and injury, it does disregard the idea that the weakest sarcomeres lengthen first as suggested by the sarcomere popping theory.<sup>10</sup>

An alternative explanation is the winding filament theory, which proposes titin binds to actin upon the presence of calcium, in turn reducing titin's "free length".<sup>13</sup> Therefore, the strain placed on titin, and its contribution to force, is greater for an absolute increase in length. In the absence of calcium (passive stretch), titin does not bind to actin and has a greater free length.<sup>13</sup> Thus, whilst the increased force from titin during active contractions may be useful in an isolated event to prevent further lengthening, it may also accumulate muscle damage due to the excessive lengthening of the unbound or free part of titin. Furthermore, after a bout of eccentric exercise the content of titin in a myofibril is depleted, most likely due to direct damage to titin and removal by proteolysis.<sup>14</sup> Considering titin also helps stabilise myosin in

a sarcomere,<sup>15</sup> any damage to titin would also potentially leave myosin vulnerable to deformation and possibly injury. Indeed, it has been demonstrated that maximally stimulated fibres require just a 30% increase in length for injury to occur whereas passively stretched fibres can reach up to a 50% increase in length before injury.<sup>16</sup>

Greater fibre strain is observed in contractions initiated at longer lengths compared to those initiated at shorter lengths ( $p < 0.002$ ),<sup>17</sup> which is believed to increase injury risk. However, we propose the strain placed on titin should also be considered which might be greater in contractions initiated at shorter muscle lengths. Indeed, when the absolute lengthening is equal between groups, the stimulation of a muscle preceding lengthening, results in greater reductions in peak torque after a bout of eccentric exercise ( $p = 0.003$ ),<sup>17</sup> suggesting greater muscle damage. In theory, the stimulation of a muscle preceding contraction would bind titin to actin earlier, placing more strain on titin during lengthening and may explain the greater muscle damage. Although further reductions in peak torque are observed in contractions initiated at longer lengths, this is likely due to the absolute muscle length also being greater. This also explains why peak torque during an exercise bout is larger in contractions initiated at long compared to short muscle lengths ( $p > 0.05$ ).<sup>17</sup>

It has been observed that sprinters suffer injuries to the bicep femoris long head (BF) (active contraction),<sup>18</sup> whereas dancers, performing movements such as the splits,<sup>19</sup> often suffer injuries to the semimembranosus (SM) (passive contraction).<sup>20</sup> In passive conditions, the BF may have greater extensibility leaving the SM vulnerable to injury, but in active contractions, the BF may become the shortest due to the shortening of titin when titin binds to actin. Perhaps the extensibility of titin under active conditions is a risk factor for hamstring strain injuries. Indeed reduced active flexibility appears to be a risk factor for hamstring strain injuries<sup>21</sup> which is in contrast to passive measures of flexibility.<sup>22</sup> Interestingly, the length of the muscle (fascicle length) would not differ between active and passive contractions and so a

variable that is mediated by active contractions, such as titin, is likely responsible for this finding. Furthermore, it appears that the isoform of titin alters and becomes stiffer with age, resulting in larger passive tension ( $p < 0.05$ ), independent of collagen.<sup>23</sup> This may be part of the reasons older athletes are more susceptible to hamstring strain injuries.<sup>24</sup>

Continuing with the specific case of the hamstrings, isokinetic testing of hamstring eccentric strength is a weak risk factor for injury (odds ratio= 1.37;  $p = 0.04$ ),<sup>25</sup> yet strength measured during the Nordic curl (an eccentric movement) is a much stronger risk factor (relative risk= 2.9;  $p = 0.01$ ).<sup>26</sup> It would be logical to assume that isokinetic tests may be a more specific test of hamstring injury risk due to the longer muscle lengths used (hip in 90° flexion and knee achieving full extension)<sup>25</sup> compared to the Nordic curl (typically 50° joint amplitude with hip in neutral)<sup>27</sup> yet this is not the case. Tests at short muscle lengths may be a measure of eccentric strength from the active component of the sarcomere, which may be more meaningful for injury prevention. At long muscle lengths, the role of structural components such as titin may increase torque production.<sup>13</sup> Future research should determine whether passive tension, for which titin is believed to be a large contributor, in the hamstrings is a risk factor for injury. In essence, strain injuries could be dependent on not only the strain experienced by the muscle but also the contraction type and the length contraction is initiated from.

It is acknowledged this would be a complicated hypothesis to test directly with respect to occurrence of hamstring strain injuries. However, we would expect to observe greater muscle damage in the hamstrings from contractions that are initiated at shorter muscle lengths than contractions that are initiated at longer lengths providing the absolute lengthening of the muscle is equal. Furthermore, it would be expected that strength towards the end of the range of motion would be greater in contractions initiated at short lengths possibly due to titin's greater contribution.

**Authors' contributions**

CP conceived the idea and wrote the initial draft. JS and KN provided feedback and assisted with the completion of the manuscript.

**Competing interests**

None of the authors declare competing financial interests.

Accepted Manuscript

## References:

1. McCully K, Faulkner J. Injury to skeletal muscle fibers of mice following lengthening contractions. *J Appl Physiol* (1985) 1985;**59**:119-26.
2. Faulkner JA, Jones DA, Round JM. Injury to skeletal muscles of mice by forced lengthening during contractions. *Q J Exp Physiol* 1989;**74**:661-70.
3. Van Hooren B, Bosch F. Is there really an eccentric action of the hamstrings during the swing phase of high-speed running? part I: a critical review of the literature. *J Sport Sci* 2016;12:1-9.
4. Herzog W. Eccentric vs. concentric muscle contraction: that is the question. *J Sport Health Sci* 2017. doi: <http://dx.doi.org/10.1016/j.jshs.2017.01.006>
5. Yu B, Liu Hu, Garrett WE. Mechanism of hamstring muscle strain injury in sprinting. *J Sport Health Sci* 2017. <http://dx.doi.org/10.1016/j.jshs.2017.02.002>
6. Liu Y, Sun Y, Zhu W, Yu J. The late swing and early stance of sprinting are most hazardous for hamstring injuries. *J Sport Health Sci* 2017. <http://dx.doi.org/10.1016/j.jshs.2017.01.011>
7. Yu B, Liu Hu, Garrett WE. Comment on “The late swing and early stance of sprinting are most hazardous for the hamstring injuries” by Liu et al. *J Sport Health Sci* 2017. <http://dx.doi.org/10.1016/j.jshs.2017.02.003>
8. Liu Y, Sun Y, Zhu W, Yu J. Comments to “Mechanism of hamstring muscle strain injury in sprinting” by Yu et al. *J Sport Health Sci* 2017. <http://dx.doi.org/10.1016/j.jshs.2017.01.012>
9. Li L, Wang D. Parallel and cross sectional hamstring injuries in sprint running. *J Sport Health Sci* 2017. <http://dx.doi.org/10.1016/j.jshs.2017.03.002>
10. Morgan DL, Proske U. Popping sarcomere hypothesis explains stretch induced muscle damage. *Clin Exp Pharmacol Physiol* 2004;**31**:541-5.

11. Morgan DL. New insights into the behaviour of muscle during active lengthening. *Biophys J* 1990;**57**:209-21.
12. Johnston K, Jinha A, Herzog W. The role of sarcomere length non-uniformities in residual force enhancement of skeletal muscle myofibrils. *R Soc Open Sci* 2016;**3**:150657. doi: 10.1098/rsos.150657. eCollection 2016
13. Leonard TR, Herzog W. Regulation of muscle force in the absence of actin-myosin-based cross-bridge interaction. *Am J Physiol Cell Physiol* 2010;**299**:C14-20.
14. Trappe TA, Carrithers JA, White F, Lambert CP, Evans WJ, Dennis RA. Titin and nebulin content in human skeletal muscle following eccentric resistance exercise. *Muscle Nerve* 2002;**25**:289-92.
15. Herzog W. The role of titin in eccentric muscle contraction. *J Exp Biol* 2014;**217**:2825-33.
16. Brooks SV, Zebra E, Faulkner JA. Injury to muscle fibres after single stretches of passive and maximally stimulated muscles in mice. *J Physiol* 1995;**488**: 459-69.
17. Butterfield TA, Herzog W. Effect of altering starting length and activation timing of muscle on fiber strain and muscle damage. *J Appl Physiol (1985)* 2006;**100**:1489-98.
18. Askling CM, Tengvar M, Saartok T, Thorstensson A. Acute first-time hamstring strains during high-speed running: a longitudinal study including clinical and magnetic resonance imaging findings. *Am J Spor Med* 2007;**35**:197-206.
19. Askling C, Saartok T, Thorstensson A. Type of acute hamstring strain affects flexibility, strength, and time to return to pre-injury level. *Br J Sports Med* 2006;**40**:40-4.
20. Askling CM, Tengvar M, Saartok T, Thorstensson A. Acute first-time hamstring strains during slow-speed stretching: clinical, magnetic resonance imaging, and recovery characteristics. *Am J Sports Med* 2007;**35**:1716-24.



21. Henderson G, Barnes CA, Portas MD. Factors associated with increased propensity for hamstring injury in English Premier League soccer players. *J Sci Med Sport* 2010;13:397-402.
22. Yeung SS, Suen AM, Yeung EW. A prospective cohort study of hamstring injuries in competitive sprinters: preseason muscle imbalance as a possible risk factor. *Br J Sports Med* 2009;43:589-94.
23. Ottenheijm CA, Knottnerus AM, Buck D, Luo X, Greer K, Hoying A, et al. Tuning passive mechanics through differential splicing of titin during skeletal muscle development. *Biophys J* 2009;97:2277-86.
24. Gabbe BJ, Finch CF, Bennell KL, Wajswelner H. Risk factors for hamstring injuries in community level Australian football. *Br J Sports Med* 2005;39:106-10.
25. van Dyk N, Bahr R, Whiteley R, Tol JL, Kumar BD, Hamilton B, et al. Hamstring and quadriceps isokinetic strength deficits are weak risk factors for hamstring strain injuries: a 4-year cohort study. *Am J Sports Med* 2016;44:1789-95.
26. Timmins R, Bourne M, Shield A, Williams M, Opar D.. Strength and architectural risk factors for hamstring strain injury in elite Australian soccer: a prospective cohort study. *Br J Sports Med* 2015;50:1524-35
27. Ditroilo M, De Vito G, Delahunt E. Kinematic and electromyographic analysis of the nordic hamstring exercise. *J Electromyogr Kinesiol* 2013;23:1111-8.